Chemistry 3A - Spring 2001 Midterm 1

Professor	Jean	Fréchet
1 10103301	ocan	LICOLICE

February 20, 2001

/ou	themistry 3A_Frechet_	
Prir	spring 2001_midterm 1	

You

Please check the section number and name of your GSI/1A.

161	Padilla-De Jesus, Omayra	371	Miljanic, Ognjen
171	Fox,Daniel	311	Sivamani,Raja
181	Furuta, Paul	321	Li, Ben
191	Ling,Frank	461	Huang,Alan
111	Cordaro, Joseph	×471	Liang, Catherine
121	Le,Scheherazade	411	Phillips,Scott
131	Thalji,Reema	421	Saxon, Eliana
141	Catherine Seeley	431	Osterhout, Robin
261	Peterka, Darcy	561	Merolle, Mauro
271	Miljanic, Ognjen	511	Klopp, John
211	Dertz, Emily	521	Wu,Sarah
221	Simon, Matthew	531	Rao, Vikas
361	Barry.David		

If you are making up an I-grade, indicate the semester you took 3A____ and the Professor .

This exam has 10 pages; make sure that you have them all.

Please be sure to use the <u>very</u> useful data given on page 10.

We will only grade answers that are in the designated spaces. Please do your scratch work on the backs of the exam pages. Write only **one** answer to each problem; multiple answers will receive **no** credit, even if one of them is correct.

Note: This examination runs for a total of 90 minutes. No questions will be answered by proctors after the exam begins. Please write legibly; ambiguous or messy answers will receive no credit.

2 b, C Zeroney 4ii (Grau) 8 (Davuol)


d

© 2001 J.M.J. Fréchet, Berkeley, CA

1. (10 points) Name or draw, as appropriate, the following molecules.

a. Br 3 a

(use IUPAC rules of nomenclature)




(use IUPAC rules of nomenclature)

c. Cis-t-butyl-4-methylcyclohexane
(Most stable conformation
use the template provided)

d. Trans-1-cyclopropyl-2-methylcyclopentane

e. 7,7-dichloro-2-methylspiro[5.2]octane

2. (16 points) (a) Draw the two chair conformations of trans-1,4-dimethylcyclohexane.



<u>--</u>

Calculate ΔG^{o} , explaining the origin of the energy difference.

Calculate the	ratio of the two	conformers at	100°C	show all w	ork

ratio =

(b) Draw a structure for each of the following molecules in the more stable chair conformation.

3. (12 points)	
3a. Draw a sawhorse projection of the least stable conformation of hexane CH ₃ (CH ₂) ₄ CH ₃ as seen along the C ₃ -C ₄ bond.	3b Draw a Newman projection of the GAUCHE conformation of butane C ₄ H ₁₀ seen along the C ₂ -C ₃ bond.
3b. Write a balanced chemical equation for the combust carbon dioxide is produced by the combustion of 44gran	ion of propane C ₃ H ₈ and calculate how much ns of propane (use C = 12, H = 1, O = 16).
balanced chemical equation	
	Weight of carbon = dioxide produced
3c. Match the name of a famous chemist with the fol Kekule, Bohr, Planck, Einstein, Pasteur, van der Waal	lowing (choose amongst: Grignard, Lewis, s, Pauling, Newman, Pauli, Shrodinger).
(i) "No orbital may be occupied by more than two electrons"	
(ii) The concept of intermolecular attractive forces	

4. (13 points)

(i) Which of the following is the stronger acid, (a) water or (b) ammonia?

Answer:

(ii) 2,4-pentanedione (pKa = 9) is a stronger acid than 2-propanone (pKa = 22). Note that in each structure below the proton involved in acid base chemistry is highlighted. Write the structure for the conjugate base of each acid and consider possible resonance forms, then explain why 2,4-pentanedione is more acidic than 2 propanone.

 $H_3C-C-CH-C-CH_3$ 2,4-pentanedione O H

H₃C-C-CH₂-H 2-propanone

iii) Write the electronic configuration and show the orbital energy diagram for Al13,




5. (12 points).

(a) Name the scientist who introduced the following concept: "Bonds are made by the in-phase overlap of atomic orbitals".

Answer:

(b) Write the Lewis-dot structure for nitrogen N₂ and for nitric acid HNO₃ (HONO₂). (Show charges if any).





(c) What is the shape and approximate H-C-C bond angle for the molecule of ethyne C_2H_2 ? Draw it in Kekule form and explain your answer using VSEPR.



- (d) What does the abbreviation VSEPR stand for?
- (e) Using arrows, show which atoms of the molecule of trimethylamine oxide have a formal charge. Specify the formal charge by writing it clearly (e.g.: +3) next to the arrow

$$H_3C$$
 N CH_3 CH_3

6. (14 points)

(a) What is the hybridization of C in ethene C₂H₄. Draw the orbitals involved in bonding the two C atoms to each other. Label these orbitals showing their type (e.g. s or sp, etc..), Explain what types of bonds are formed between the two carbons and indicate their type with a Greek letter.

(b) What is the approximate bond angle in the following triatomic molecules?

BeH₂

Answer:

H2S

Answer:

(c) Name the process used to convert alkanes into aromatic compounds.

Answer:

(d) Arrange the following alkanes in order of increasing boiling points.

- (1) 2-methylbutane (2) propane
- (3) pentane (4) 2,2-dimethylpropane

highest lowest

7. (11 Points

(a) In 1,2-dichloroethane at room temperature, the ratio of molecules in the ANTI to GAUCHE conformation is 7.6 to 1. Calculate the difference in energy between these two conformations. Show equations and detailed calculations.

	-/	 -			
Answer:					

(b) Calculate the equilibrium constant for the reaction below.

CH₃COOH + NH₃ CH₃COO + NH₄ Show detailed equations & calculations.

Answer: Keq =

8. (12 points) (a) The monobromination of propane affords one major product. Write the propagation steps for the preparation of the major product and calculate ΔH^0 for the overall reaction.



(b) Draw and label the energy diagram for the formation of the major monobromination product of propane.

Note: There are no questions to be answered on this page, it only contains data that may be of use in solving the questions contained in this exam. Not all of the data given is needed.

Value of gas constant: R = 2.0 cal deg-1 mol-1

 pK_a of $CH_3COOH = 4.76$ pK_a of $NH_4^+ = 9.24$

Value of e (base for natural logarithms) e = 2.718

Value of absolute zero (kelvin) = -273°C

Bond dissociation energies (in kcal mole⁻¹): RCH₂-H 98 R₂CH-H 95 R₃C-H 91; CI-CI 58; Br-Br 46; H-Br 88; H-CI 103 RCH₂-CI 81

RCH2-Br 68 R2CH-Cl 80 R2CH-Br 68 R3C-Cl 79; R3C-Br 65;

Values of strain energies:

Each CH₃ - H eclipsing interaction: 1.5 kcal mol⁻¹

Each H - H eclipsing interaction: 1.0 kcal mol-1

Each CH₃ - CH₃ eclipsing interaction: 2.5 kcal mol⁻¹

Each CH₃ - CH₃ butane-gauche interaction: 0.9 kcal mol⁻¹

Each t-Butyl - CH₃ gauche interaction: 2.0 kcal mol⁻¹

Each CH₃ - H 1,3-diaxial interaction: 0.9 kcal mol⁻¹

Each CI - H 1,3-diaxial interaction: 0.25 kcal mol-1

Each CH₃ - CH₃ 1,3-diaxial interaction: 1.6 kcal mol⁻¹

Each H - CN 1,3-diaxial interaction: 0.1 kcal mol-1

Each H - C(CH₃)₃ 1,3-diaxial interaction: 2.5 kcal mol⁻¹

Partial periodic table of the elements

1 H 1.00794	IIA			IIIA	IVA	VA	VIA	VIIA	O He 4.00260
3 Li 6.941	Be 9.01218			5 B 10.811	6 C 12.011	7 N 14.0067	8 0 15.9994	9 F 18.9984	10 Ne 20.1797
11 Na 22.9898	12 Mg 24.3050	IB	IIB	13 Al 26.9815	14 Si 28.0855	15 P 30.9738	16 S 32.066	17 CI 35.4527	18 Ar 39.948
19 K 39.0983	20 Ca 40.078	 29 Cu 63.546	30 Zn 65.39	31 Ga 69.723	32 Ge 72.61	33 As 74.9216	34 Se 78.96	35 Br 79.904	36 Kr 83.80