

University of California, Berkeley Physics

137A Spring 2011

MidTerm Exam I

Maximum score: 100 points

1. (50 points)

Considering the one-dimensional problem of a particle of mass  $m$  in a potential

$$V = 0 \quad x < -a$$

$$V = -\frac{40\hbar^2}{ma^2}, \quad -a < x < 0$$

$$V = \infty \quad x > 0$$

- (1) Sketch the potential and the first 3 energy eigenfunctions.
- (2) Solve symbolically the energy eigenvalues and eigenfunctions.
- (3) How many bound states exist in this potential?
- (4) Now considering a scattering state coming from  $-\infty$  with energy of  $\frac{\hbar^2}{ma^2}$ . What will the reflectivity?

2. (50 points)

A Schrodinger equation in one dimension reads  $-\partial^2\psi/\partial x^2 - 2\text{sech}^2x \cdot \psi = E\psi$ . (We have set  $\hbar=1$  and  $m=1/2$ .)

- (1) Show that  $\exp(ikx)(\tanh x + \overset{\text{constant}}{\downarrow})$  is a solution of the Schrodinger equation for a particular value of constant. Calculate the transmission and reflection coefficients.
- (2) Using analytic extension of the transmission function of (1) to find the eigenenergy and eigenfunction of one bound state in this potential.
- (3) Sketch the wavefunction you found in (2). Give a simple argument that it must be the ground state of the potential.

Definition:  $\text{sech}(x) \equiv \frac{2}{e^x + e^{-x}}, \tanh(x) \equiv \frac{e^{2x} - 1}{e^{2x} + 1}$ .