
Physics 7b Spring 2000 Midterm 2 R. Packard

Sharm & Solution 1/5


Name:	
SID	
GSI name	Section number (or time)
the question ask the proctor for clarifi	ghted equally. If you don't understand cation. Do not perform any numerical raic answer. Do a dimension check on
$e=1.6x10^{-19}C$ , $N_A=6.02x10^{23}$ , $k_B=1.38$	$8 \times 10^{-23} \text{JK}^{-1}$ , $\varepsilon_0 = 8.85 \times 10^{-12}$
1	
2	
3	
4	
Total	

1. A point charge +Q is located at the origin and is surrounded by a spherical charge distribution that varies proportional to the distance from the origin:  $\rho=\alpha r$  for  $r \le R$ . The charge density drops to zero when r>R. The total charge is zero. A) Find the constant  $\alpha$ . B) Find the electric field for all r.

2. A uniformly charged ring of radius a and charge  $Q_1$  lies in the yz plane with its axis along the x axis. A point charge  $Q_2$  is placed on the x axis at x=2a. Find the work required to move the point charge to the origin.



- 3. A) Find the capacitance of the parallel plate capacitor shown in the figure. The plate separation is 1mm, the plate area is  $25 \times 10^{-2} \text{m}^2$ ,  $k_1=2$ ,  $k_2=4$  and  $k_3=3$ . The dielectric slabs on the left are of equal thickness and take up half the area between the plates.
- B) Assume a capacitor with similar plate area and separation is filled with a single dielectric slab of dielectric constant k=3. This capacitor is charged to 200V. How much work is required to pull the dielectric completely out of the capacitor. Neglect friction.



4. A copper wire (mass density  $\rho_m=8.9x10^3$  kg/m³,  $M_w=63x10^{-3}$ kg, conductivity  $\sigma=6x10^7$  ( $\Omega$ m)<sup>-1</sup>) is 100m long and has a diameter of 0.8mm. Under the influence of a current an electron takes  $3x10^5$ s to travel the length of the wire. What is the electric field in the wire?